

AMENDMENTS TO THE CLAIMS

1. (Currently amended) A method for producing a synchronized population of conifer somatic embryos, the method comprising:

(a) cultivating pre-cotyledonary conifer embryogenic cells in, or on a maintenance medium comprising nutrients that sustain the embryos and one or more agents for adjusting the osmolality of the medium to a desired range;

~~[[a)]~~ (b) cultivating the pre-cotyledonary conifer embryogenic cells from step (a) for a period of at least 0.5 week in, or on, a synchronization medium that comprises an absorbent composition and at least one synchronization agent selected from the group consisting of abscisic acid and a gibberellin, wherein the absorbent composition and the at least one synchronization agent are present at a concentration effective to produce a synchronized population of pre-cotyledonary conifer somatic embryos wherein at least 50% of the embryos in the synchronized population are at the same developmental stage; and

~~[[b)]~~ (c) transferring the synchronized population of pre-cotyledonary conifer somatic embryos from step (b) to a development medium for synchronized cotyledonary embryo development.

2. (Original) The method of Claim 1 wherein the absorbent composition is selected from the group consisting of activated charcoal, soluble poly(vinyl pyrrolidone), insoluble poly(vinyl pyrrolidone), activated alumina, and silica gel.

3. (Original) The method of Claim 2 wherein the absorbent composition is activated charcoal.

4. (Original) The method of Claim 1 wherein the concentration of the absorbent composition in the synchronization medium is from about 0.5 g/L to about 50 g/L.

LAW OFFICES OF
CHRISTENSEN O'CONNOR JOHNSON KINDNESS^{PLLC}
1420 Fifth Avenue
Suite 2800
Seattle, Washington 98101
206 682 8100

5. (Original) The method of Claim 1 wherein the absorbent composition is activated charcoal, and the activated charcoal is present in the synchronization medium at a concentration in the range of from about 0.1 g/L to about 5 g/L.

6. (Original) The method of Claim 1 wherein the absorbent composition is activated charcoal, and the activated charcoal is present in the synchronization medium at a concentration in the range of from about 0.5 g/L to about 1 g/L.

7. (Original) The method of Claim 1, wherein abscisic acid is used as a synchronization agent.

8. (Original) The method of Claim 1, wherein a gibberellin is used as a synchronization agent.

9. (Original) The method of Claim 1, wherein abscisic acid and at least one gibberellin are used as synchronization agents.

10. (Original) The method of Claim 1, wherein a gibberellin is present in the synchronization medium at a concentration of from about 0.5 mg/L to about 500 mg/L.

11. (Original) The method of Claim 1, wherein a gibberellin is present in the synchronization medium at a concentration of from about 1.0 mg/L to about 100 mg/L.

12. (Original) The method of Claim 1, wherein abscisic acid is present in the synchronization medium at a concentration of from about 1.0 mg/L to about 500 mg/L.

13. (Original) The method of Claim 1, wherein abscisic acid is present in the synchronization medium at a concentration of from about 0.5 mg/L to about 20 mg/L.

14. (Previously presented) The method of Claim 1, wherein the conifer embryogenic cells are cultured in, or on, the synchronization medium for a period of from about 0.5 week to about 5 weeks.

15. (Original) The method of Claim 1, wherein the conifer embryogenic cells are cultured in, or on, the synchronization medium for a period of from about 1 week to about 3 weeks.

16. (Original) The method of Claim 1, wherein the conifer embryogenic cells are cultured in, or on, the synchronization medium for a period of from about 1 week to about 2 weeks.

17. (Original) The method of Claim 1, wherein the osmolality of the synchronization medium is from about 90 mM/Kg to about 300 mM/Kg.

18. (Original) The method of Claim 1, wherein the pH of the synchronization medium is from about 5 to about 6.

19. (Original) The method of Claim 1, wherein the conifer somatic embryos are produced from Loblolly pine embryogenic cells.

20. (Canceled)

21. (Original) The method of Claim 1, wherein at least 75% of the embryos in the synchronized population of conifer somatic embryos are at the same developmental stage.

22. (Canceled)